

Exploring retropharyngeal lymph node metabarcoding as a tool for characterizing microbial signatures in white-tailed deer from urban and non-urban areas

George Zaragoza¹ , Samuel Greaves¹ , Melanie Almanza¹, Breityn Barnhart¹, Caitlin Blackburn¹, Aidan Breen¹ , Jessica Dorrien¹, Michael David Garan¹ , Jenay Lemasney¹, Lee Nguyen¹, Katlene Patron¹, Bruno Vinicius Pinheiro¹, Bianca Pinilla¹, Ellie Rowland¹, Bambi Clemons², & Meguine Duvert-Ortega²

Collaborators: Tyler Brasington, Grace Kahler + 2 other reviewers

Accepted by 4 of 4 reviewers

Conflicts of Interest

The authors declare no conflicts of interest.

Publishing History

Submitted March 5 2026
Accepted September 9 2026
Published October 29 2026

Corresponding Author

George Zaragoza
george.zaragoza@ucf.edu



Open Access



Peer-Reviewed



Creative Commons

¹ Department of Biology, University of Central Florida, Orlando, Florida, USA

² Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission, Gainesville, Florida, USA

Disease Ecology

Genomics

Mammals

Monitoring

Urban Ecology

Abstract

Evaluating the effects of urbanization on wildlife has become increasingly important for understanding how host-associated microbial communities respond to environmental change. Characterizing these communities can improve our understanding



of shifts in disease pressure and provide information for early detection of changes. Wildlife disease screening programs may also provide a source of opportunistically collected samples for research and monitoring. Here, we evaluated the use of retropharyngeal lymph nodes from white-tailed deer (*Odocoileus virginianus*) collected in Florida as part of chronic wasting disease surveillance to assess bacterial community composition across urban and non-urban areas. We detected no differences in alpha or beta microbial diversity between deer from urban versus non-urban environments. However, we found evidence of isolation-by-distance in beta diversity, with communities becoming more dissimilar as geographic distance increased and localized clustering near some urban areas. This study explores how lymph nodes collected during routine surveillance can be used to monitor spatial variation and temporal changes in host-associated microbial communities, and we provide recommendations for researchers interested in applying and improving this approach in future studies.

Keywords: diversity, disease, pathogen, monitoring, Florida

Introduction

Wildlife living in urban environments are exposed to a variety of novel stressors that can affect the establishment and composition of species communities (Faeth et al., 2011). For example, urban environments may expose wildlife to increased levels of chemical pollutants, higher temperatures, and pathogens (Bradley and Altizer, 2007, Sepp et al., 2019; Swank et al., 2021). Human land-use modification has also previously been associated with changes to disease prevalence, including chronic wasting disease (CWD) in mule deer (Farnsworth et al., 2005). In gut microbiomes, resident microbial communities can influence pathogen colonization through competition for resources, inhibitions, and interactions with host immune responses (Pickard et al., 2017). In white ibises, urban-associated shifts in gut microbiome composition have also coincided with lower



microbial diversity, and individuals with lower diversity were more likely to shed *Salmonella* (Murray et al., 2020). Evaluating the differences in microbial community composition across urban and non-urban areas can therefore aid in identifying shifts associated with urbanization and potentially pathogenic taxa of concern for disease surveillance.

White-tailed deer (*Odocoileus virginianus*) are one species commonly found in both urban and non-urban areas of Florida. White-tailed deer in urban environments have demonstrated changes in behavior compared to their non-urban counterparts, with higher likelihood of temporal and spatial overlap with human activity (Péter et al., 2025). Urban deer can have overlapping home ranges, potentially increasing the rate of disease transmission (Walter et al., 2011). Additionally, simulations have suggested that fragmented landscapes can concentrate white-tailed deer within remaining habitat fragments and accelerate the transmission of CWD (Kjaer and Schaubert, 2022), which has also been positively correlated with human land-use modification in mule deer (Farnsworth et al., 2005). Although previous research in *Odocoileus* has found limited differences in body condition and pathogen prevalence between urban and non-urban individuals (Mathieu et al., 2018), few studies have leveraged microbial community characterization across urban gradients to establish baseline community profiles and monitor shifts in microbial communities, including the detection of potentially pathogenic taxa that may require more focused disease surveillance. This monitoring can support more responsive pathogen-specific screening and also contribute to One Health initiatives focused on mitigating spillover events (Flint et al., 2015, Eby et al., 2023, Hayman et al., 2023).

Evaluation of bacterial community composition and diversity is often done through methods such as metabarcoding, which enables low-cost and rapid screening of diverse sample types with varying quality (Taberlet et al., 2012, Cabodevilla et al., 2024). This flexibility can be particularly useful for analyzing opportunistically collected tissues such as lymph nodes, which have previously been used to assess microbial community composition in harvested wildlife and commercial livestock, and to monitor food sources for pathogens prior to human consumption (Wittekindt et al., 2010, Mann et al., 2015, Queirós et al., 2019). Lymph nodes are also the primary sites of infection by various pathogens, including several *Salmonella* species (St



John and Abraham, 2009). Because lymph nodes process lymph draining from tissues, microbial signatures may reflect exposure to local environmental factors (Hirosue and Dubrot, 2015). This makes them particularly valuable for wildlife and disease monitoring programs, such as CWD surveillance, where they are regularly collected and stored.

Here, we assessed microbial community structure in lymph nodes obtained from white-tailed deer in Florida. We performed 16S metabarcoding of individuals collected across urban and non-urban areas and assessed differences in community diversity across urban and non-urban groups. We found geographically correlated patterns of diversity between locations and identified taxa that may be of interest for disease management and monitoring. We also provide suggestions for improving the methodology of future studies that leverage lymph node sample collections for research in microbial community assessment or surveillance.

Methods and Materials

Sample collection & DNA extraction

We obtained retropharyngeal lymph nodes from 77 *O. virginianus* sampled by the Florida Fish and Wildlife Conservation Commission (FWC) as part of CWD screening in the state. These samples included individuals that were primarily identified as roadkill and hunter harvested, as well as those that died of other causes between the years 2019 and 2021, with lymph nodes later harvested and frozen. To assign individuals to urban or nonurban areas, we used the United States Census Bureau's 2020 Urban Area shapefile, which delineates urban and metropolitan areas with high population densities and urban land use (U.S. Census Bureau, 2022).

We generated a buffer with a 1 km radius, equivalent to a home range area of approximately 3.14 km², around each occurrence using the Buffer geoprocessing tool in QGIS v3.40.14 (QGIS Development Team, 2026). We chose this buffer as its area falls within the range of published white-tailed deer home range estimates in Florida, which include a mean adult female range of 1.94 km² and mean annual adult male home range of 4.54-7.0 km²



(Land et al., 1993, Sargent and Labisky, 1995). We calculated the proportion of urban land within each buffer based on its intersection with the U.S. Census Bureau urban area polygon (Figure 1).




Figure 1. Sampling locations of white-tailed deer in central and southern Florida. Pie charts show the proportion of urban (purple) and non-urban (green) land cover within a 1 km radius buffer around each sampling location. The locations of overlapping pies have been slightly adjusted for visualization.



Library preparation and sequencing

We extracted DNA via spin-column using a Quick-DNA Miniprep Plus Kit (Zymo Inc., Irvine, California, USA) per the manufacturer's instructions, using an overnight digestion of a lymph node fragment with Proteinase K at 55°C. We targeted the V3-V4 hypervariable regions of the 16S rRNA following the Illumina 16S Metagenomic Sequencing Library Preparation workflow. We amplified genomic DNA in 25 µL PCR reactions containing 12.5µL KAPA HiFi HotStart ReadyMix (Kapa Biosystems Inc., Wilmington, Massachusetts, USA), 5 µL of forward and reverse primers with Illumina overhang adapters (forward: 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGA CAGCCTACGGGNGGCWGCAG-3'; reverse: 5'-GTCTCGTGGGCTCGGAGATGT GTATAAGAGACAGGACTACHVGGGTATCTAATCC-3') and 2.5 µL of DNA extract. We performed initial PCR under the following conditions: 90 °C for 3 min, 25 cycles of 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s, and final extension at 72 °C for 5 min. PCR was followed by magnetic bead cleanups to remove primers and short fragments.

Sequencing libraries were created by attaching Illumina sequencing adapters and unique dual indices through indexing PCR. Reactions contained 25 µL KAPA HiFi ReadyMix, 10 µL water, 10 µL IDT for Illumina unique dual indices (Illumina Inc., San Diego, California, USA), and 5 µL of purified PCR product. Indexing PCR was run under the following conditions: 95 °C for 30 s, 55 °C for 30 s, and 72 °C for 30 s, and final extension at 72 °C for 5 min. We also purified indexed libraries with a second magnetic bead cleanup to remove residual adapters and primers.

We quantified final libraries using qPCR with the NEBNext Library Quant Kit for Illumina. We prepared serial dilutions of each library and quantified them using manufacturer-provided DNA standards using Illumina adapter-specific primers. We performed qPCR under the following conditions: initial denaturation at 95 °C for 1 min, 35 cycles of 95 °C for 15 s, and 63 °C for 45 s. We performed equimolar pooling of libraries prior to sequencing. Sequencing was performed on an Illumina MiSeq platform (v3 chemistry, 2 x 300 bp paired-end reads) at the University of Florida Interdisciplinary Center for Biotechnology Research (ICBR) NextGen DNA Sequencing Core.



Bioinformatic analyses

We processed raw FASTQ files using QIIME2 v2025.10 (Bolyen et al., 2019) and assessed sequencing quality and read depth across samples. Forward and reverse primer sequences and adapters were removed with qiime cutadapt trim-paired plugin (Martin, 2011). Amplicon sequence variants (ASVs) were inferred using the DADA2 algorithm in qiime dada2 denoise-paired (Callahan et al., 2016). We truncated forward and reverse reads to 234 bp and 229 bp, respectively, based on quality score distributions while maintaining a median quality score of 30 across the read length (Supplementary Material). We then filtered and merged paired reads using qiime dada2 denoise-paired to generate a ASV feature table and representative ASV sequences. We identified and removed contaminants using qiime quality-control decontam-identify, using negative controls defined in sample metadata. Contaminant filtering was performed using a decontam score threshold of 0.10, and negative controls were removed from the decontaminated feature table.

We assigned taxonomy to ASVs using the SILVA ribosomal RNA reference database (Chuvpochina et al., 2025) with qiime feature-classifier classify-sklearn. We retained taxonomic assignments at the highest confidence supported by the classifier and filtered the feature table to remove sequences annotated as mitochondria or chloroplasts, restricting analyses to bacterial taxa. We also reconstructed phylogenetic relationships in ASVs prior to diversity analyses by aligning representative sequencing using MAFFT (Kato and Standley, 2013) and constructing a phylogenetic tree using FastTree (Price et al., 2010). We generated rarefaction curves using qiime diversity alpha-rarefaction and selected a rarefaction cutoff of 1000 sequences per sample to balance sample retention and stabilization of diversity estimates across samples (Supplementary Material).

We evaluated alpha microbial diversity using the Shannon diversity index and observed features (ASV richness) and beta microbial diversity using Bray-Curtis dissimilarity and Jaccard distance across all rarefied samples. Significance of associations between alpha diversity and the proportion of urban land within the 1 km radius buffers was tested using Spearman correlations. For beta diversity, we evaluated associations based on the



proportion of urban land using permutation multivariate analysis of variance (PERMANOVA) (Anderson, 2017) with 999 permutations. We also tested whether alpha diversity and beta diversity differed by sex and age class using Kruskal-Wallis tests. To assess spatial structure and isolation-by-distance, we generated pairwise geographic distance matrices from sample coordinates and performed Mantel tests to test correlations between geographic distance and community dissimilarity for Bray-Curtis and Jaccard distance matrices.

Results

The 77 sampled deer included 48 males and 29 females, which were primarily adults (N = 45), with vehicle collisions (N = 41) and hunter harvests (N = 18) as the primary causes of death (Table 1). We generated 7,250,147 paired-end read pairs across 77 samples after adapter trimming, averaging 94,158 read pairs per sample. Following DADA2 quality filtering, paired-end merging, and removal of contaminant reads and negative controls, 5,353,462 reads remained across 72 samples, representing 3,805 ASVs with lengths ranging from 234-451 nucleotides. Resulting sample sequencing depth had a median frequency of 8,177 (IQR 1348-33,666). Of the 3,805 ASVs, 2,709 remained after removal of mitochondrial and chloroplast sequences. At the phylum level, the taxa with the highest prevalence across the 72 samples were Firmicutes (N = 70), Proteobacteria (N = 70), Actinobacteriota (N = 53), Bacteroidota (N = 49), and Patescibacteria (N = 21). At the genus level, the taxa with the highest prevalence included *Pseudomonas* (N = 61), *Psychobacter* (N = 50), *Escherichia-Shigella* (N = 50), *Streptococcus* (N = 47), *Acinetobacter* (N = 42), *Cutibacterium* (N = 40), *Vagococcus* (N = 33), *Clostridium sensu stricto* (N = 30), *Staphylococcus* (N = 28), and *Carnobacterium* (N = 26). Other notable taxa that were present included *Erysipelothrix* (N = 23) and *Bartonella* (N = 20).



Table 1. Sex, cause of death, and age classes of samples white-tailed deer grouped by urban classification, with urban samples described as those having at least 50% urban land within the 1 km radius buffer around collection locations.

		Overall (N=77)	Non-Urban (<50% urban, N=48)	Urban (≥50% urban; N=29)
Sex	Male	48	36	12
	Female	29	21	8
Cause of death	Vehicle	41	27	14
	Hunting	18	17	1
	Fence entanglement	7	6	1
	Predation	1	0	1
	Unknown	10	7	3
Age category	Adult	45	34	11
	Yearling	23	19	4
	Fawn	8	4	4
	Unknown	1	0	1

After rarefaction to 1000 sequences per sample, 50 (69.4%) of the 72 samples were retained, including 23 from urban areas and 27 from non-urban areas. The mean number of observed features (ASV richness) was 41.5 ($\sigma = 24.8$) for urban samples and 53.6 ($\sigma = 46.1$) for non-urban samples, and there was no evidence of differentiation between groups ($H = 0.27$, $p = 0.61$). Mean Shannon diversity was 3.47 ($\sigma = 1.23$) for urban and 3.19 ($\sigma = 1.54$) for non-urban, with no differences between groups ($H = 0.72$, $p = 0.40$). No patterns or clustering associated with the proportion of urban land or other metadata were apparent in Bray-Curtis or Jaccard ordinations (Supplementary Material). PERMANOVA showed no significant association between the proportion of urban land and community composition for Bray-Curtis ($F = 0.73$, $R^2 = 0.01$, $p = 0.89$) and Jaccard ($F = 1.12$, $R^2 = 0.02$, $p = 0.11$). Additionally, we did not detect significant differences in alpha or beta diversity by sex (Bray-Curtis: $F = 1.01$, $R^2 = 0.02$, $p = 0.42$; Jaccard: $F = 0.90$, $R^2 = 0.2$, $p = 0.87$) or age class (Bray-Curtis: $F = 0.68$, $R^2 = 0.02$, $p = 0.99$; Jaccard: $F = 0.96$, $R^2 = 0.03$, $p = 0.74$). Results of our Mantel test revealed significant associations between geographic distance and community dissimilarity based on both Bray-Curtis ($r = 0.26$, $p < 0.01$) and Jaccard ($r = 0.25$, $p < 0.01$) beta diversity, and there was some geographical clustering based on principal component ordination, particularly near the Tampa metropolitan urban area (Figure 2).




Figure 2. Geographic distribution of variation in bacterial community composition in white-tailed deer samples (following rarefaction) from central and southern Florida. Sampling locations are colored by scores along the first principle coordinate axis (PCoA1) from Jaccard dissimilarities (A) and Bray-Curtis dissimilarities (B). Samples with similar colors have more similar bacterial community composition, whereas contrasting colors indicate greater differences. The locations of overlapping points have been slightly adjusted for visualization.

Discussion

We used lymph nodes collected from CWD monitoring to evaluate differences in community composition in deer from urban and non-urban areas in Florida. Although we were unable to detect differentiation using alpha and beta diversity statistics, we found that patterns of beta diversity were geographically correlated using the Bray-Curtis and Jaccard indices. These results suggest that urbanization alone, when represented using broadly defined urban areas, may not be the only driver of lymph node-associated microbial community diversity across our sampled deer, and that the observed variation may be due to other processes, including more localized spatial or individual-level processes. The apparent geographical clustering near the Tampa metropolitan area, but lack of significant differentiation in diversity statistics, suggests that bacterial diversity may be driven more by a specific urban area's population



dynamics than by general urban versus non-urban patterns. Habitat fragmentation can concentrate deer within habitat patches and increase overlap among individuals (Water et al., 2011, Kjaer and Schaubert, 2022), and we may not be able to detect variation in small urban habitat patches or at finer spatial scales where interactions might take place. Additionally, local differences in deer density, movement, and contact rates can also contribute to the observed clustering independently of, or in addition to, the proportion of urban land cover. We classified several taxa at the genus level, and the most common across samples were *Pseudomonas*, *Psychrobacter*, *Escherichia-Shigella*, *Streptococcus*, and *Acinetobacter*. *Pseudomonas*, *Psychrobacter*, and *Acinetobacter* are broad genera that include both pathogenic and non-pathogenic environmental and host-associated taxa (Wong et al., 2016, Welter et al., 2021, Girard et al., 2021). *Escherichia-Shigella* is a potential pathogen commonly found in gut microbiomes (Li et al., 2018, 2022), and its detection in lymph node tissue may reflect transfer during handling of deer carcasses rather than a lymph node-specific signal. This may be particularly relevant for deer collected after a vehicle collision, which may remain at the site of mortality for an unknown period prior to collection and have greater environmental exposure and leakage of bodily fluids. *Streptococcus* is common in vertebrate oral and upper respiratory microbiomes, but the genus also encompasses both commensal and pathogenic lineages (Abranches et al., 2018). Other taxa with potential relevance to disease monitoring included *Erysipelothrix* and *Bartonella*. *Erysipelothrix* includes the ungulate pathogen *Erysipelothrix rhusiopathiae*, which has been associated with widespread mortality in northern ungulates (Forde et al., 2016, Mavrot et al., 2020). *Bartonella* is a bloodborne vector-transmitted genus frequently detected in cervids and includes zoonotic species that can cause bacteremia in humans (Duodu et al., 2013, Razanske et al., 2018). Although these results provide a descriptive overview of retropharyngeal lymph node-associated bacterial taxa, interpretation of the function or impact of these taxa on communities of white-tailed deer should be approached cautiously without species- or strain-specific identification.

Disease monitoring by wildlife agencies can provide an opportunistic source of retropharyngeal lymph node samples and similar tissues, and we offer four recommendations for researchers interested in leveraging these for surveillance and monitoring. First, metabarcoding using multiple



markers can broaden detection and improve taxonomic resolution. Standard 16S rRNA metabarcoding of the V3-V4 hypervariable region primarily targets bacterial DNA and is not intended to detect fungi, eukaryotic microorganisms, or viruses (Bukin et al., 2019). We recommend including additional metabarcoding targets such as fungal ITS (Beeck et al., 2014) and eukaryotic 18S rRNA (Hadziavdic et al., 2014) regions, as well as viral metabarcoding or metagenomic approaches (Gil et al., 2021). Second, QIIME databases such as SILVA are useful for broad taxonomic profiling, though we encourage researchers to develop custom or disease-specific databases to provide more relevant characterization for monitoring beyond a general community overview. These databases typically include curated sequences with taxonomic annotations and can be tailored to detect species or taxa of particular concern, for example through tools such as RESCRIPt, which support the development and creation of custom reference databases (Robeson et al., 2021). Third, sample collection and storage may be a major source of variation, especially for hunter-harvested samples and field collection with resource limitations. When possible, we recommend preservation by freezing or ethanol storage to mitigate post-mortem bacterial growth. We also encourage researchers to partner with agencies to engage hunters as collaborators by communicating the value of these samples for disease monitoring, which can promote more consistent handling, confidence in results, and help support healthy game populations through proactive disease management. Fourth, future work should focus on sampling multiple habitat patches within specific urban areas where managers or researchers have knowledge of local areas that support deer, to improve detection of patterns that may be missed with broad characterizations of urbanization. Habitat patches within urban and suburban areas can form mosaics that create complex dynamics relevant to potential disease transmission. Expanding the number of sampling sites in urban areas would improve characterization of microbial communities, particularly as deer can occur within major cities where suitable habitat patches remain (Ciach and Fröhlich, 2019, Potratz et al., 2019).

Conclusion

In this study, we characterized bacterial communities in retropharyngeal lymph nodes collected through CWD surveillance to evaluate patterns



associated with urbanization and geography in Florida deer. We found no differences between urban and non-urban groups in alpha or beta diversity. However, geographic structuring in beta diversity suggests local spatial patterns. These results highlight the utility of wildlife disease surveillance programs, such as CWD screening, as an opportunistic source of tissues for research and monitoring beyond their original intent. Using these tissues for long-term monitoring may also provide a proactive means of detecting shifts in diversity associated with changing disease risk or pressure as part of an integrated wildlife monitoring approach (Barroso et al., 2023). We recommend that researchers use multi-marker approaches to broaden detection across taxa, develop disease monitoring-relevant databases for characterization, preserve samples as consistently as feasible to minimize post-mortem changes, and prioritize sampling in specific urban centers to better understand local patterns of diversity.

Acknowledgments

We thank Chris Sarkis for the maintenance and care of laboratory equipment and samples used in this study. We thank Dr. Anna Savage and Dr. Anna Forsman for providing protocols and recommendations for metabarcoding library preparation. We thank Shouq Alajmi and Colin Shoening for their contributions to the laboratory work and initial interpretation and reporting of the data. This work was conducted as part of an undergraduate research intensive genomics course offered by the Department of Biology at the University of Central Florida. We thank the Biology department for funds that supported this work.

Funding Information

There is no funding to report for this project.

Author Contributions

G. Zaragoza: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project Administration, Supervision, Writing – original draft, review & editing



S. Greaves: Formal analysis, Investigation, Methodology, Project Administration, Supervision, Writing – original draft, review & editing
M. Almanza: Investigation, Writing – original draft, review & editing
B. Barnhart: Investigation, Writing – original draft, review & editing
C. Blackburn: Investigation, Writing – original draft, review & editing
A. Breen: Investigation, Writing – original draft, review & editing
J. Dorrien: Investigation, Writing – original draft, review & editing
M. D. Garan: Investigation, Writing – original draft, review & editing
J. Lemasney: Investigation, Writing – original draft, review & editing
L. Nguyen: Investigation, Writing – original draft, review & editing
K. Patron: Investigation, Writing – original draft, review & editing
B. V. Pinheiro: Investigation, Writing – original draft, review & editing
B. Pinilla: Investigation, Writing – original draft, review & editing
E. Rowland: Investigation, Writing – original draft, review & editing
B. Clemons: Data curation, Methodology, Writing – review & editing
M. Duvet-Ortega: Data curation, Methodology, Writing – review & editing

Data Availability

Sequencing reads generated in this study are available in the NCBI Sequence Read Archive under BioProject PRJNA1419337 at <https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1419337>. Scripts used to perform the data analyses in this paper are available at <https://github.com/georgeomics>.

Supplementary Information

Supplementary information [can be found here](#).

Transparent Peer Review

Results from the Transparent Peer Review [can be found here](#).



Recommended Citation

Zaragoza, G., S. Greaves, M. Almanza, B. Barnhart, C. Blackburn, A. Breen, J. Dorrien, M. D. Garan, J. Lemasney, L. Nguyen, K. Patron, B. Vinicius Pinheiro, B. Pinilla, E. Rowland, B. Clemons, and M. Duvet-Ortega. 2026. Exploring retropharyngeal lymph node metabarcoding as a tool for characterizing microbial signatures in white-tailed deer from urban and non-urban areas. *Stacks Journal*: 26013. <https://doi.org/10.60102/stacks-26013>.



References

- Abranches, J., L. Zeng, J. K. Kajfasz, S. R. Palmer, B. Chakraborty, Z. T. Wen, V. P. Richards, L. J. Brady, J. A. Lemos. 2018. Biology of oral Streptococci. *Microbiology Spectrum* 6. <https://doi.org/10.1128/microbiolspec.GPP3-0042-2018>.
- Anderson, M. J. 2006. Distance-based tests for homogeneity of multivariate dispersions. *Biometrics* 62: 245-253. <https://doi.org/10.1111/j.1541-0420.2005.00440.x>.
- Anderson, M. J. 2017. Permutational multivariate analysis of variance (PERMANOVA). *in: Wiley StatsRef: Statistics Reference Online*, John Wiley & Sons, Ltd. 1-15. <https://doi.org/10.1002/9781118445112.stat07841>.
- Barroso P., D. Relimpio, J. A. Zearra, J. J. Cerón, P. Palencia, B. Cardoso, E. Ferreras, M. Escobar, G. Caceres, J.R. López-Olvera, and C. Gortázar. 2023. Using integrated wildlife monitoring to prevent future pandemics through one health approach. *One Health* 16: 100479. <https://doi.org/10.1016/j.onehlt.2022.100479>.
- Beaumelle, C., E. M. Redman, J. De Rijke, J. Wit, S. Benabed, F. Debias, J. Duhayer, S. Pardonnet, M.T. Poiriel, G. Capron, and S. Chabot. 2021. Metabarcoding in two isolated populations of wild roe deer (*Capreolus capreolus*) reveals variation in gastrointestinal nematode community composition between regions and among age classes. *Parasites Vectors* 14: 594. <https://doi.org/10.1186/s13071-021-05087-5>.
- Bolyen, E., J. R. Rideout, M. R. Dillon, N. A. Bokulich, C. C. Abnet, G. A. Al-Ghalith, et al. 2019. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology* 37: 852-857. <https://doi.org/10.1038/s41587-019-0209-9>.
- Bradley, C. A., and S. Altizer 2007. Urbanization and the ecology of wildlife diseases. *Trends in Ecology & Evolution* 22: 95-102. <https://doi.org/10.1016/j.tree.2006.11.001>.
- Bukin, Y. S., Y. P. Galachyants, I. V. Morozov, S. V. Bukin, A. S. Zakharenko, and T. I. Zemskaya. 2019. The effect of 16S rRNA region choice on bacterial community metabarcoding results. *Scientific Data* 6: 190007. <https://doi.org/10.1038/sdata.2019.7>.
- Cabodevilla, X., J. E. Malo, D. Aguirre de Cárcer, J. Zurdo, R. Chaboy-Cansado, A. Rastrojo, et al. 2024. Zoonotic potential of urban wildlife faeces, assessed through metabarcoding. *Science of The Total Environment* 952: 175866. <https://doi.org/10.1016/j.scitotenv.2024.175866>.
- Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016. DADA2: high-resolution sample inference from Illumina amplicon data. *Nature Methods* 13: 581-583. <https://doi.org/10.1038/nmeth.3869>.
- Chuvochina, M., J. Gerken, M. Frentrup, Y. Sandikci, R. Goldmann, H. M. Freese, et al. 2025. SILVA in 2026: a global core biodata resource for rRNA within the DSMZ digital diversity. *Nucleic Acids Resources: gkaf1247*. <https://doi.org/10.1093/nar/gkaf1247>.
- Ciach, M., and A. Fröhlich 2019. Ungulates in the city: light pollution and open habitats predict the probability of roe deer occurring in an urban environment. *Urban Ecosystems* 22: 513-523. <https://doi.org/10.1007/s11252-019-00840-2>.
- De Beeck M. O., B. Lievens, P. Busschaert, S. Declerck, J. Vangronsveld, and J. V. Colpaert. 2014. Comparison and validation of some ITS primer pairs useful for fungal metabarcoding studies. *PLOS ONE* 9: e97629. <https://doi.org/10.1371/journal.pone.0097629>.
- Duodu, S., K. Madslien, E. Hjelm, Y. Molin, A. Paziewska-Harris, P. D. Harris, et al. 2013. *Bartonella* infections in deer keds (*Lipoptena cervi*) and moose (*Alces alces*) in Norway. *Applied and Environmental Microbiology* 79: 322-327. <https://doi.org/10.1128/AEM.02632-12>.



- Eby, P., A. J. Peel, A. Hoegh, W. Madden, J. R. Giles, P. J. Hudson, et al. 2023. Pathogen spillover driven by rapid changes in bat ecology. *Nature* 613: 340-344. <https://doi.org/10.1038/s41586-022-05506-2>.
- Faeth, S. H., C. Bang, and S. Saari. 2011. Urban biodiversity: patterns and mechanisms. *Annals of the New York Academy of Sciences* 1223: 69-81. <https://doi.org/10.1111/j.1749-6632.2010.05925.x>.
- Farnsworth, M. L., L. L. Wolfe, N. T. Hobbs, K. P. Burnham, E. S. Williams, D. M. Theobald, et al. 2005. Human land use influences chronic wasting disease prevalence in mule deer. *Ecological Applications* 15: 119-126. <https://doi.org/10.1890/04-0194>.
- Flint, P. L., J. M. Pearce, J. C. Franson, and D. V. Derksen. 2015. Wild bird surveillance for highly pathogenic avian influenza H5 in North America. *Virology Journal* 12: 151. <https://doi.org/10.1186/s12985-015-0377-2>.
- Forde, T. L., K. Orsel, R. N. Zadoks, R. Biek, L. G. Adams, S. L. Checkley, et al. 2016. Bacterial genomics reveal the complex epidemiology of an emerging pathogen in Arctic and boreal ungulates. *Frontiers in Microbiology* 7. <https://doi.org/10.3389/fmicb.2016.01759>.
- Gil, P., V. Dupuy, R. Koual, A. Exbrayat, E. Loire, A. G. Fall, et al. 2021. A library preparation optimized for metagenomics of RNA viruses. *Molecular Ecology Resources* 21: 1788-1807. <https://doi.org/10.1111/1755-0998.13378>.
- Girard, L., C. Lood, M. Höfte, P. Vandamme, H. Rokni-Zadeh, V. van Noort, et al. 2021. The ever-expanding *Pseudomonas* genus: description of 43 new species and partition of the *Pseudomonas putida* group. *Microorganisms* 9: 1766. <https://doi.org/10.3390/microorganisms9081766>.
- Hadziavdic, K., K. Lekang, A. Lanzen, I. Jonassen, E. M. Thompson, and C. Troedsson. 2014. Characterization of the 18S rRNA gene for designing universal eukaryote specific primers. *PLOS ONE* 9: e87624. <https://doi.org/10.1371/journal.pone.0087624>.
- Hayman, D. T. S., W. B. Adisasmito, S. Almuhairi, C. B. Behraves, P. Bilivogui, S. A. Bukachi, et al. 2023. Developing One Health surveillance systems. *One Health* 17: 100617. <https://doi.org/10.1016/j.onehlt.2023.100617>.
- Hirosue, S., and J. Dubrot. 2015. Modes of antigen presentation by lymph node stromal cells and their immunological implications. *Frontiers in Immunology* 6. <https://doi.org/10.3389/fimmu.2015.00446>.
- Katoh, K., and D. M. Standley. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772-780. <https://doi.org/10.1093/molbev/mst010>.
- Kjær, L. J., and E. M. Schaubert. 2022. The effect of landscape, transmission mode and social behavior on disease transmission: simulating the transmission of chronic wasting disease in white-tailed deer (*Odocoileus virginianus*) populations using a spatially explicit agent-based model. *Ecological Modelling* 472: 110114. <https://doi.org/10.1016/j.ecolmodel.2022.110114>.
- Klindworth, A., E. Pruesse, T. Schweer, J. Peplies, C. Quast, M. Horn, et al. 2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* 41: e1. <https://doi.org/10.1093/nar/gks808>.
- Land, E. D., D. S. Maehr, J. C. Roof, and J. W. McCown. 1993. Mortality patterns of female white-tailed deer in southwest Florida. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 47: 176-184.
- Lewis, J. S., K. A. Logan, M. W. Alldredge, S. Carver, S. N. Bevins, M. Lappin, et al. 2017. The effects of demographic, social, and environmental characteristics on pathogen prevalence in wild felids across a gradient of urbanization. *PLOS ONE* 12: e0187035. <https://doi.org/10.1371/journal.pone.0187035>.



- Li, S., J. Guo, R. Liu, F. Zhang, S. Wen, Y. Liu, et al. 2022. Predominance of *Escherichia-Shigella* in gut microbiome and its potential correlation with elevated level of plasma tumor necrosis factor alpha in patients with tuberculous meningitis. *Microbiology Spectrum* 10: e01926-22. <https://doi.org/10.1128/spectrum.01926-22>.
- Li, Z., X. Wang, T. Zhang, H. Si, W. Nan, C. Xu, et al. 2018. The development of microbiota and metabolome in small intestine of sika deer (*Cervus nippon*) from birth to weaning. *Frontiers in Microbiology* 9. <https://doi.org/10.3389/fmicb.2018.00004>.
- Mann, E., B. Pinior, S. U. Wetzels, B. U. Metzler-Zebeli, M. Wagner, and S. Schmitz-Esser. 2015. The metabolically active bacterial microbiome of tonsils and mandibular lymph nodes of slaughter pigs. *Frontiers in Microbiology* 6. <https://doi.org/10.3389/fmicb.2015.01362>.
- Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* 17: 10-12. <https://doi.org/10.14806/ej.17.1.200>.
- Mathieu, A., M. Flint, P. M. Stent, H. M. Schwantje, and T. E. Wittum. 2018. Comparative health assessment of urban and non-urban free-ranging mule deer (*Odocoileus hemionus*) in southeastern British Columbia, Canada. *PeerJ* 6: e4968. <https://doi.org/10.7717/peerj.4968>.
- Mavrot, F., K. Orsel, W. Hutchins, L. G. Adams, K. Beckmen, J. E. Blake, et al. 2020. Novel insights into serodiagnosis and epidemiology of *Erysipelothrix rhusiopathiae*, a newly recognized pathogen in muskoxen (*Ovibos moschatus*). *PLOS ONE* 15: e0231724. <https://doi.org/10.1371/journal.pone.0231724>.
- Murray, M. H., E. W. Lankau, A. D. Kidd, C. N. Welch, T. Ellison, H. C. Adams, et al. 2020. Gut microbiome shifts with urbanization and potentially facilitates a zoonotic pathogen in a wading bird. *PLOS ONE* 15: e0220926. <https://doi.org/10.1371/journal.pone.0220926>.
- Péter, S. A., T. Gallo, J. Mullinax, A. Roess, G. Palomo-Munoz, and T. Anderson. 2025. Integrating human mobility and animal movement data reveals complex space-use between humans and white-tailed deer in urban environments. *Scientific Reports* 15: 18588. <https://doi.org/10.1038/s41598-025-03577-5>.
- Pickard, J. M., M. Y. Zeng, R. Caruso, and G. Núñez. 2017. Gut microbiota: role in pathogen colonization, immune responses, and inflammatory disease. *Immunological Reviews* 279: 70-89. <https://doi.org/10.1111/imr.12567>.
- Potratz, E. J., J. S. Brown, T. Gallo, C. Anchor, and R. M. Santymire. 2019. Effects of demography and urbanization on stress and body condition in urban white-tailed deer. *Urban Ecosystems* 22: 807-816. <https://doi.org/10.1007/s11252-019-00856-8>.
- Price, M. N., P. S. Dehal, and A. P. Arkin. 2010. FastTree 2 – approximately maximum-likelihood trees for large alignments. *PLOS ONE* 5: e9490. <https://doi.org/10.1371/journal.pone.0009490>.
- QGIS Development Team. 2026. QGIS geographic information system. QGIS Association. <https://www.qgis.org>.
- Queirós, J., M. Villar, A. Hernández-Jarguín, V. López, I. Fernández de Mera, J. Vicente, et al. 2019. A metaproteomics approach reveals changes in mandibular lymph node microbiota of wild boar naturally exposed to an increasing trend of *Mycobacterium tuberculosis* complex infection. *Tuberculosis* 114: 103-112. <https://doi.org/10.1016/j.tube.2018.12.003>.
- Razanske, I., O. Rosef, J. Radzijeuskaja, K. Klepeckiene, I. Lipatova, and A. Paulauskas. 2018. Infections with *Bartonella* spp. in free-ranging cervids and deer keds (*Lipoptena cervi*) in Norway. *Comparative Immunology, Microbiology and Infectious Diseases* 58: 26-30. <https://doi.org/10.1016/j.cimid.2018.06.003>.



- Robeson, M. S., D. R. O'Rourke, B. D. Kaehler, M. Ziemski, M. R. Dillon, J. T. Foster, et al. 2021. RESCRIPT: reproducible sequence taxonomy reference database management. *PLOS Computational Biology* 17: e1009581. <https://doi.org/10.1371/journal.pcbi.1009581>.
- Sargent, R. A., and R. F. Labisky. 1995. Home range of male white-tailed deer in hunted and non-hunted populations. *Proceedings of the Annual Conference of the Southeastern Association of Fish and Wildlife Agencies* 49: 389-398.
- Sepp, T., B. Ujvari, P. W. Ewald, F. Thomas, and M. Giraudeau. 2019. Urban environment and cancer in wildlife: available evidence and future research avenues. *Proceedings of the Royal Society B* 286: 20182434. <https://doi.org/10.1098/rspb.2018.2434>.
- St. John, A. L., and S. N. Abraham. 2009. *Salmonella* disrupts lymph node architecture by TLR4-mediated suppression of homeostatic chemokines. *Nature Medicine* 15: 1259-1265. <https://doi.org/10.1038/nm.2036>.
- Swank, A., L. Wang, J. Ward, and H. Schoenfuss. 2021. Multigenerational effects of a complex urban contaminant mixture on the behavior of larval and adult fish in multiple fitness contexts. *Science of The Total Environment* 791: 148095. <https://doi.org/10.1016/j.scitotenv.2021.148095>.
- Taberlet, P., E. Coissac, F. Pompanon, C. Brochmann, and E. Willerslev. 2012. Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* 21: 2045-2050. <https://doi.org/10.1111/j.1365-294X.2012.05470.x>.
- U.S. Census Bureau (2022). TIGER/Line shapefile, 2020, nation, U.S., 2020 census urban area.
- Walter, W. D., J. Beringer, L. P. Hansen, J. W. Fischer, J. J. Millspaugh, and K. C. Vercauteren. 2011. Factors affecting space use overlap by white-tailed deer in an urban landscape. *International Journal of Geographical Information Science* 25: 379-392. <https://doi.org/10.1080/13658816.2010.524163>.
- Welter, D. K., A. Ruaud, Z. M. Henseler, H. N. De Jong, P. van Coeverden de Groot, J. Michaux, et al. 2021. Free-living, psychrotrophic bacteria of the genus *Psychrobacter* are descendants of pathobionts. *mSystems* 6. <https://doi.org/10.1128/msystems.00258-21>.
- Wittekindt, N.E., A. Padhi, S. C. Schuster, J. Qi, F. Zhao, L. P. Tomsho, L. R. Kasson, M. Packard, P. Cross, and M. Poss. 2010. Nodeomics: Pathogen detection in vertebrate lymph nodes using meta-transcriptomics. *PLOS ONE* 5: e13432. <https://doi.org/10.1371/journal.pone.0013432>.
- Wong, D., T. B. Nielsen, R. A. Bonomo, P. Pantapalangkoor, B. Luna, and B. Spellberg. 2016. Clinical and pathophysiological overview of *Acinetobacter* infections: a Century of challenges. *Clinical Microbiology Reviews* 30: 409-447. <https://doi.org/10.1128/CMR.00058-16>.